



## DEPARTMENT OF MATHEMATICS

### UNIT-I PARTIAL DIFFERENTIAL EQUATIONS

#### HOMOGENEOUS EQUATIONS:

$$(D^2 + 2DD' + D'^2)z = \cos(x-y)$$

A.E. is  $m^2 + 2m + 1 = 0$

$\Rightarrow m = -1, m = -1$

C.F. is  $z = f_1(y-x) + \alpha f_2(y-x)$

P.I. =  $\frac{1}{D^2 + 2DD' + D'^2} \cos(x-y)$

$\frac{1}{-1+2-1} \cos(x-y)$

D.w.r. to 'D' in 'Dr' & multi. by x in the 'Nr'.

=  $x \frac{1}{2D + 2D'} x \cos(x-y)$

=  $\frac{1}{2D + 2D'} \times \frac{2D - 2D'}{2D - 2D'} x \cos(x-y)$

=  $\frac{2x[D - D'] \cos(x-y)}{4D^2 - 4D'^2}$

$\begin{cases} D \rightarrow \text{D.w.r. to } x \\ D' \rightarrow \text{D.w.r. to } y \end{cases}$

$D^2 = -(1)^2 = -1$

$D'^2 = -(-1)^2 = -1$

=  $2x [D(\cos(x-y)) - D'(\cos(x-y))]$

=  $4(-1) = 4(-1)$



## DEPARTMENT OF MATHEMATICS

### UNIT-I PARTIAL DIFFERENTIAL EQUATIONS

$$[D^2 - D D' - 2D'^2]z = x^2y + (2x + 3y)$$

$$\text{AE: } m^2 - m - 2 = 0 \Rightarrow m = 2, -1$$

$$\text{CF: } z = f_1(y-x) + f_2(y+2x)$$

$$\begin{aligned} \text{P.I.:} &= \frac{1}{D^2 - D D' - 2D'^2} x^2y \\ &= \frac{1}{D^2 \left[ 1 - \frac{(D D' + 2D'^2)}{D^2} \right]} x^2y = \frac{1}{D^2} \left[ 1 - \left( \frac{D'}{D} + \frac{2D'^2}{D^2} \right) \right]^{-1} x^2y \end{aligned}$$

$$= \frac{1}{D^2} \left[ 1 + \frac{D'}{D} + \frac{2D'^2}{D^2} \right] x^2y$$

$$= \frac{1}{D^2} \left[ x^2y + \frac{D'}{D} (x^2y) + \frac{2D'^2}{D^2} (x^2y) \right]$$

$$= \frac{1}{D^2} \left[ x^2y + \frac{1}{D} \frac{d}{dy} (x^2y) + \frac{2}{D^2} \frac{d^2}{dy^2} (x^2y) \right]$$

$$= \frac{1}{D^2} \left[ x^2y + \frac{1}{D} (x^2) + \frac{2}{D^2} (0) \right]$$

$$= \frac{1}{D^2} \left[ x^2y + \frac{x^3}{3} \right]$$

$$= \frac{1}{D} \int \left( x^2y + \frac{x^3}{3} \right) dx$$

$$= \int \left( \frac{x^3}{3} y + \frac{x^4}{12} \right) dx$$

$$= \frac{x^4}{12} y + \frac{x^5}{60}$$



# SNS COLLEGE OF TECHNOLOGY

(An Autonomous Institution)

Coimbatore – 35



## DEPARTMENT OF MATHEMATICS

### UNIT-I PARTIAL DIFFERENTIAL EQUATIONS

$$\begin{aligned}
 P.I_2 &= \frac{1}{D^2 - DD' - 2D'^2} (2x+3y) \\
 &= \frac{1}{D^2 \left[ 1 - \left( \frac{D'}{D} + \frac{2D'^2}{D^2} \right) \right]} (2x+3y) \\
 &= \frac{1}{D^2} \left[ 1 - \left( \frac{D'}{D} + \frac{2D'^2}{D^2} \right) \right]^{-1} (2x+3y) \\
 &= \frac{1}{D^2} \left[ 1 + \frac{D'}{D} + \frac{2D'^2}{D^2} \right] (2x+3y) \\
 &= \frac{1}{D^2} \left[ 2x+3y + \frac{D'}{D} (2x+3y) + \frac{2D'^2}{D^2} (2x+3y) \right]
 \end{aligned}$$

**TYPE IV :** RHS =  $f(my) = e^{ax+by} x^m y^n$  (or)  $e^{ax+by} \cos(ax+by)$  or  $\sin(ax+by)$

$$P.I = \frac{1}{\phi(D, D')} e^{ax+by} x^m y^n.$$

Replace  $D \rightarrow D+a$  ;  $D' \rightarrow D+b$ . Then type III rule or type II rule.



## DEPARTMENT OF MATHEMATICS

### UNIT-I PARTIAL DIFFERENTIAL EQUATIONS

1) Solve:  $(D^2 - 2DD' + D'^2)z = x^2y^2 \cdot e^{x+y}$

Soln: A.E. is  $m^2 - 2m + 1 = 0$

$$(m-1)^2 = 0$$

$$m = +1, +1$$

$\therefore$  The roots are real and equal.

C.F. is  $z = f_1(y+mx) + x f_2(y+mx)$   
 $= f_1(y+x) + x f_2(y+x)$

P.I.  $P.I. = \frac{1}{D^2 - 2DD' + D'^2} \cdot x^2y^2 e^{x+y}$

Replace  $D \rightarrow D+1$ ;  $D' \rightarrow D'+1$

$$= \frac{1}{(D+1)^2 - 2(D+1)(D'+1) + (D'+1)^2} \cdot e^{x+y} \cdot x^2y^2$$



## DEPARTMENT OF MATHEMATICS

### UNIT-I PARTIAL DIFFERENTIAL EQUATIONS

$$= \frac{1}{D^2 + 2D + 1 - 2[D\partial' + D + \partial' + 1]} + D'^2 + 2D' + 1 \cdot e^{x+y} \cdot x^2 y^2$$

$$= \frac{1}{D^2 + 2D + 1 - 2D\partial' - 2D - 2\partial' - 2} + D'^2 + 2D' + 1 \cdot e^{x+y} \cdot x^2 y^2$$

$$= \frac{1}{D^2 - 2D\partial' + D'^2} \cdot e^{x+y} \cdot x^2 y^2$$

$$= e^{x+y} \cdot \frac{1}{D^2} \left[ 1 - \left( \frac{2D\partial'}{D} - \frac{D'^2}{D^2} \right) \right]^{-1} x^2 y^2$$

$$= e^{x+y} \cdot \frac{1}{D^2} \left[ 1 + \left( \frac{2D\partial'}{D} - \frac{D'^2}{D^2} \right) \right] x^2 y^2$$

$$= e^{x+y} \cdot \frac{1}{D^2} \left[ x^2 y^2 + \frac{2D\partial'}{D} x^2 y^2 - \frac{D'^2}{D^2} x^2 y^2 \right]$$

$$= e^{x+y} \cdot \left[ \frac{1}{D^2} (x^2 y^2) + \frac{2}{D^2} \left( 2 \frac{x^3 y}{3} \right) - \frac{1}{D^2} \left( 2 \frac{x^4}{12} \right) \right]$$

$$= e^{x+y} \cdot \left[ \frac{x^4 y^2}{12} + \frac{4y}{3} \frac{x^5}{20} - \frac{1}{6} \frac{x^6}{30} \right]$$

$$= e^{x+y} \cdot \left[ \frac{x^4 y^2}{12} + \frac{x^5 y}{15} - \frac{x^6}{180} \right]$$

∴ Solution is  $z = C.F + P.I$

$$= f_1(y+x) + x f_2(y+x) + e^{x+y} \left[ \frac{x^4 y^2}{12} + \frac{x^5 y}{15} - \frac{x^6}{180} \right]$$